

Utilization of Sewage Effluents for Irrigating Crop Plants for Sustainable Improvement I – Percentage Seed Germination and Seed Germination Index in Cereal and Legume Crops

Snehlata¹, Nidhi Parashar (Sharma)², Abha Pathak³, Kapila Saraswat³ and Ravi Sharma^{4*}

¹Research Scholar Department of Botany and Scientific Assistant UP Pollution Control Board Regional Office Mathura

²Research Scholar Department of Chemistry

³Research Scholars Department of Botany

⁴Formerly Head Department of Botany, K.R. College, Mathura and Ex-Founder Principal ESS College of Education Dayalbagh Agra (Dr B. R. Ambedkar Univ., Agra Formerly Agra University, Agra) UP India

***Corresponding Author:** Ravi Sharma, Formerly Head Department of Botany, K.R. College, Mathura and Ex-Founder Principal ESS College of Education Dayalbagh Agra (Dr B. R. Ambedkar Univ., Agra Formerly Agra University, Agra) UP India.

Received: October 01, 2017; **Published:** May 23, 2018

Abstract

The present study included physico-chemical analysis of Sewage Effluents (SE) entering the River Yamuna at Mathura UP India with effect of SE water on leguminous and cereal crops of the region observed to assess its potential utilization for irrigating crop plants for increasing improved and sustainable agricultural production. As such SE was analyzed and its effect in various concentrations (25%, 50%, 75% and 100% along with controls as distilled water) on seed germination was performed on two cereal (wheat and barley) and two legume (mustard and broad bean) crops. Mustard and Broad bean did not show any seed germination at all in all the SE treatment concentrations at 24 hours duration. Besides, in Broad bean no germination was observed at 48 hours too in all the SE treatments. Further, it was observed that 75% SE treatment and also 72 hours of germination were found to be quite critical for all the crops. The results in relation to the effect of different concentrations of SE exposure on germination performance measured in terms of Germination Percentage (GP) and Speed of Germination Index (SGI) have shown germination performance as C3 Mustard > C4 Broad bean > C1 Wheat > C2 Barley at critical duration and critical level of 72 hours in 75% SE concentration showing better performance of Mustard and Broad bean than Wheat and Barley.

Keywords: Sewage Effluent (SE); Germination Percentage (GP); Speed of Germination Index (SGI); Cereal Crops; Legume Crops;

Volume 2 Issue 6 May 2018

© All Copy Rights are Reserved by Ravi Sharma., *et al.*

The steady increase in industrialization, urbanization and enormous population growth are leading to production of huge quantities of wastewaters that may frequently cause environmental hazards. Paucity of quality fresh water for agriculture has made wastewater application a popular option. This makes wastewater treatment and wastewater reduction very important issues [Khurana and Singh, 2012; Snehlata, 2016]. The present study offers a collection of studies and findings concerning wastewater treatment, minimization and reuse.

Citation: Ravi Sharma., *et al.* "Utilization of Sewage Effluents for Irrigating Crop Plants for Sustainable Improvement I – Percentage Seed Germination and Seed Germination Index in Cereal and Legume Crops". *Innovative Techniques in Agriculture* 2.6 (2018): 487-508.

Water supply and water quality degradation are global concerns that had intensified with increasing water demand; for this reason, worldwide marginal-quality water has become an increasingly important component of agricultural water supplies particularly in water scarce regions. The status of severe water resources shortage determines that new water source must be developed to cope with the deficiency of water sources for agriculture irrigation. One of the major types of marginal-quality water is the wastewater from urban and peri-urban areas [Pedrero., *et al.* 2010]. The municipal wastewater is a potential water resource with stability of water quantity and reliable supply. Irrigation with reclaimed municipal wastewater that is properly treated and satisfied with the agricultural recycling standards has huge benefits and profound social effects [Shi., *et al.* 2008]. In recent years wastewater use has gained importance in water-scarce regions. In Pakistan 26% of vegetable production is irrigated with wastewater [Ensink., *et al.* 2004]. In Ghana, informal irrigation involving diluted wastewater from rivers and streams occurs on an estimated 11,500 hectares [Keraita., *et al.* 2014]. In Mexico about 260,000 hectares are irrigated with wastewater, mostly untreated [Mexico CAN, 2004]. In India, estimates revealed that ~73,000 hectares were irrigated with wastewater during early nineties and presently the area under wastewater irrigation is on the rise [Strauss and Blumenthal, 1990; Astera, 2014]. Wastewater reuse in agriculture is an ancient practice that has been generally implemented worldwide. Agricultural deployment of wastewater for irrigation is based on the value of its constituents, which are used as fertilizers. However, crop irrigation with insufficiently treated wastewater may result in health risks. The use of sewage effluent for irrigation exposes the public to the dangers of infection with a variety of pathogens such as bacteria, viruses, protozoa and helminths. Thus, the benefit of wastewater reuse is limited by its potential health hazards associated with the transmission of pathogenic organisms from the irrigated soil to crops, to grazing animals and humans [Gupta., *et al.* 2009]. Human health risks from wastewater irrigation include firstly farmers' and consumers' exposure to pathogens including other infections, and secondly, organic and inorganic trace elements. Protective measures such as wearing boots and gloves, and changing irrigation methods can reduce farmer exposure [Qadir, *et al.* 2010].

Further, ground water resources in most areas of world are shrinking at an alarming rate and may not meet the ever increasing demands from agriculture and industry in future. Estimates revealed that agriculture sector consumes maximum of the ground water and ~80 per cent of actual water resources are utilized in agriculture for irrigation purpose. Many farmers in areas near to urban/peri-urban localities are even compelled to use wastewater to irrigate their crops, due to absence of better alternatives [Ghimire, 1994]. Cheraghi., *et al.* [2009] revealed an annual production of ~30 million tons of wastewater in the world, of which ~70% is consumed as an agricultural fertilizer and irrigation source. Wastewater utilization for crop production has gained an acceptance all over the world [Singh., *et al.* 2004], as an economic alternate that could substitute nutrient needs [Campbell and Davidson, 1983; Burns., *et al.* 1985; Chaw and Reves, 2001; Rusan., *et al.* 2007].

However, increasing the productivity of water and making safe use of poor quality waters in agriculture will play a vital role in easing competition for scarce water resources, prevention of environmental degradation and provision of food security [Parashar, 2011; Parashar and Sharma, 2017; Parashar., *et al.* 2018]. Driven by the pressure to produce more, even the effluent rich saline and alkali waters are being increasingly diverted to irrigated agriculture [Sharma, 1982; 1987; 2016; 2017; Nauhbar, 2005; Yadav, 2007; Rani, 2008; Gautam, 2009]. There are two major approaches to improving and sustaining productivity in a poor quality water irrigation environment: modifying the environment to suit the plant and modifying the plant to suit the environment, but the former has been tried more extensively. The available options are mediated through the management of crops/sequences, irrigation water, chemical/amendments and other cultural practices but all must be integrated as per the site specific needs and achieving higher yields on sustainable basis. Some important interventions include appropriate crop/variety selection, blending effluents and fresh water to keep the resultant effluent toxicity below threshold, or their cyclic application by scheduling irrigation with salty water at less salt sensitive stages. At the farm/irrigation system level, policy measures like re-allocation of water to higher value crops and those with limited irrigation requirement, spatial re-allocation and transfer of water, adopting policies that favor development of water markets and adequate utilization of groundwater can help in improving water productivity in polluted environments. The other viable options include effluent irrigation tolerant agro-forestry systems and bio-effluent agriculture. Besides technological advances, peoples' participation and favorable water

use and allocation policies need to be put in place for sustainable use of effluent waters [Sharma, 1976; Sharma, 2004; Parashar, 2011; Snehlata, 2016; Parashar and Sharma, 2017; Parashar, *et al.* 2018].

Sewage is the water-borne waste derived from domestic waste and animal or food processing plants. It consists of human excreta and other organic wastes from kitchen and bathrooms, paper, cloth, soap, detergents etc. Raw sewage consists of about 99.9% of water and about 0.1% of solids. Seventy percent of these are organic constituents and 30% are inorganic. Sixty percent of the organic substances are proteins, while carbohydrates and fats represent 25% and 19% respectively. The inorganic fraction constitutes grit, salts and metals in varying proportions [Tebbutt, 1977]. The irrigation of grasslands and crop-fields with sewage is an ancient practice followed in Prussia for more than 450 years beginning 1559. Raw sewage has been flood irrigated in pastures at Melbourne, Australia since 1897. Mahida [1981] sites several examples where sewage has been used for irrigation. But, with the increasing urbanization and industrialization, the disposal of sewage effluents is becoming a serious problem all over the world. The composition of sewage sludge changes with the type of industries discharging their effluents into the sewer system. This material can be considered as an organic manure and soil amendment. The addition of sewage sludge to agricultural land has been reported to increase the yields of many crops [Unnamalai, *et al.* 2007; Sharma, 2004; Parashar, 2011; Snehlata, 2016; Parashar and Sharma, 2017]. Besides, many countries face periodical drought periods and a continuous deficit of good-quality fresh water resources due to severe water pollution problems. Most countries will face it within 20-25 years. Therefore, agriculture must use alternative water resources for increasing crop production for the ever growing population to feed. The basic idea is to achieve good yield of uncontaminated agricultural products as well as cash crops so as to make agriculture sustainable [Annan, 2005].

Mathura, a semi-arid region, faces water scarcity throughout the year with ground water being highly brackish and not fit for irrigation. Yamuna River and wells are the main sources of water in Mathura. There is pressing demand for supplemental water to augment crop production. Therefore, in Mathura use of sewage effluents after treatment and proper dilution with normal water making them suitable for irrigation is the most favorable approach. It serves as an additional potential source of fertilizer for agricultural use and also prevents the wastewater from being an environmental hazard in minimizing pollution load of the water-bodies [Gautam, 2009; Parashar, 2011; Snehlata, 2016; Parashar and Sharma, 2017; Parashar, 2018]. The reuse of wastewater for agricultural purposes is becoming more necessary as good quality water becomes scarce. Further, the proper use of this wastewater for irrigation has a two-fold advantage; it conserves a valuable resource and takes advantage of nutrients present in sewage effluents for crop requirements. Since the nutrient status and inferior quality of water bring major constraints in the development of agriculture, harvesting the nutrient energy of biological/organic and industrial wastewater is of prime importance for maximizing the food, feed, and fodder and fuel production [Khurana and Singh, 2012]. Thus, the wastewater can be used to irrigate the fields in rural areas and maintenance of large playgrounds, parks, nurseries and green belts in the urban areas. Besides, the wastewater irrigation may solve various associated problems of its disposal and may give a new life to the rivers, natural streams and other water-bodies and soils receiving this wastewater in which the pollution load can be minimized to a large extent [Sharma, 1976; Sharma, 2004; Parashar, 2011; Snehlata, 2016; Parashar and Sharma, 2017]. Besides, the ground water quality in Mathura region is highly saline and brackish. The rainfall is inadequate and unevenly distributed. The canal water from the lower Agra canal distribution system does not reach the remote fields of the cultivators in the villages [Gautam, 2009]. There is, therefore, a pressing demand for supplemental irrigation water to augment crop production [Parashar, 2011; Parashar, *et al.* 2016; Snehlata, 2016; Parashar and Sharma, 2017]. Thus, the present study is an effort and aims at a detailed study of potential water quality problems in the light of industrial and urban development in Mathura with an attempt to see its effects on plant growth and development especially agricultural crops of the region so that the quality of water is best utilized for irrigation to raise higher crop yields so as to make sustainable improvement of crops.

Materials and Methods

The present study was conducted at the Eco-physiology Laboratory, Department of Post-graduate Studies and Research in Botany, K R College, Mathura (Dr. B. R. Ambedkar University, formerly Agra University, Agra) UP, India.

Citation: Ravi Sharma, *et al.* "Utilization of Sewage Effluents for Irrigating Crop Plants for Sustainable Improvement I – Percentage Seed Germination and Seed Germination Index in Cereal and Legume Crops". *Innovative Techniques in Agriculture* 2.6 (2018): 487-508.

The present study includes physico-chemical analysis of sewage effluents entering the River Yamuna at Mathura. Effect of sewage on crops with sewage effluent water was observed to assess their potential utilization for irrigating crop plants for increasing improved and sustainable agricultural production. As such sewage effluent was analyzed and its effect in various concentrations (25%, 50%, 75% and 100% along with controls as distilled water) on seed germination was performed on two cereal (wheat and barley) and two legume (mustard and broad bean) crops of Mathura region.

Physico-Chemical Analysis and Characterization of the Sewage Effluents from Two Selected Areas in Yamuna River at Mathura was carried out following Johnson, *et al.* [1959], Weber [1971], Allen, *et al.* [1974], Tonapi [1980], NEERI [1981, 1988], Grob [1983], Trivedi and Goel [1986], APHA-AWWA-WPCF [1995], Manivasakam [2000], Thakur [2001], Singh [2003], Singh, *et al.* [2008], Parashar [2011], AOAC [2012], Snehlata [2016], Parashar, *et al.* [2018] for the following parameters: Water colour, odour, transparency, turbidity, temperature, pH, total alkalinity, free CO₂, DO, BOD, COD, TSS, TDS, Cl⁻, Ca²⁺, Na⁺, K⁺, Mg²⁺, SO₄²⁺, inorganic PO₄⁻, particulate PO₄⁻, total dissolved PO₄⁻, total Kjeldhal PO₄⁻, NH₃-N, NO₃-N.

The four crop varieties selected for the present study are most commonly grown in western U.P. particularly in Mathura and nearby areas. The seeds of the material involved in the present study were kindly provided by the Agriculture Research Centre Raya, Mathura UP, India.

The following crop plants were chosen for experimentation:

1. *Triticum aestivum* (Wheat) variety P.B.W. 373 family Poaceae
2. *Hordeum vulgare* (Barley) variety Barley Narendra family Poaceae
3. *Brassica campestris* (Mustard) variety Pusa Jaikisan family Brassicaceae
4. *Vicia faba* (Broad bean) local variety family Fabaceae

Studies on the Effect of Irrigation with Sewage Effluents on Germination Percentage and Speed of Germination Index

Seed lots of the four crop varieties were screened for their relative tolerance to Sewage Effluents (SE) under varying concentration levels viz., 0, 5, 50, 75 & 100 %. Distilled water was used as Control. As such seed germination and seedling growth was carried out on two cereal (wheat and barley) and two legume (mustard and broad bean) crops of the region by Garrad's Technique [Garrad, 1945], as modified by Sarin and Rao [1956], as per method of Sheoran and Garg [1978], followed by Sharma [1982, 1987, 2015, 2016, 2017] and Parashar and Sharma [2017]. Seed Germination was measured as Germination Percentage (GP) and Speed of Germination Index (SGI). The speed of germination index (SGI) was determined by following the formula of Carley and Watson [1968]: $SGI = (5 \times 1G + 4 \times 2G + 3 \times 3G + 2 \times 4G + 1 \times 5G)$ Where, 1G ——— 5G = Number of seeds germinated on the first (24 hours) to fifth (120 hours) day.

Preparation of Stock Solution

Stock solutions for irrigation with Sewage Effluents (SE) were prepared as follows:

1. 0% Solution Distilled Water (DW) was used as Controls
2. 25% Test solution was prepared with 25% SE + 75% DW
3. 50% Test solution was prepared with 50% SE + 50% DW
4. 75% Test solution was prepared with 75% SE + 25% DW
5. 100% Test solution was prepared with 100% SE

Sterilization of Seeds and HgCl₂ Salt Treatment

Three hundred seeds of each variety were surface sterilized by soaking in 0.1% (w/v) HgCl₂ for 2 min and then rinsed twice with sterile distilled water. Sterilized seeds were subjected to SE treatment in varying concentrations in petri-dishes for seed germination

containing autoclaved sterilized filter paper (Whatman No1) saturated with different concentrations of SE treatment. The petri-dishes were kept in BOD incubator at $25 \pm 2^\circ\text{C}$ temperature. Observations of the influence of SE as 25% test solution, 50% test solution, 75% test solution and 100% test solution of SE solutions and the Controls (DW grown) were recorded at 24 hour intervals from 24 hours after sowing up to the end of 120 hours. Seed germination was calculated after 24, 48, 72, 96 and 120 hours. Three replications with 100 seeds were taken for all the experimentation.

Statistical Analysis

All parameters with three replicates were analyzed by ‘Analysis of Variance’ (ANOVA) by using window SPSS package (SPSS 17.00 version). Data were expressed as the mean + standard error of the mean. Critical differences at 5 percent probability were calculated wherever the results were significant.

Results and Discussion

Statistical Analysis

All parameters with three replicates were analyzed by ‘Analysis of Variance’ (ANOVA) by using window SPSS package (SPSS 17.00 version). Data were expressed as the mean + standard error of the mean. Critical differences at 5 percent probability were calculated wherever the results were significant.

Physico-chemical Characterization of Sewage Effluents (SE) was done for Water colour, odour, transparency, turbidity, temperature, pH, total alkalinity, free CO_2 , DO, BOD, COD, TSS, TDS, Cl-, Ca^{2+} , Na^+ , K^+ , Mg^{2+} , SO_4^{2-} , inorganic PO_4^{--} , particulate PO_4^{--} , total dissolved PO_4^{--} , total Kjeldhal PO_4^{--} , $\text{NH}_3\text{-N}$, $\text{NO}_3\text{-N}$.

The results of the analysis of Sewage Effluents (SE) at Two Sites (Site I- Masani Nala and Site II-Sewage Farm) have been shown in the Table 1. Highly significant differences were noticed among all the parameters at 0.01% and 0.05% level of Probability:

S. No.	Physico-chemical Parameters	Masani Nala	Sewage Farm
1	Colour	Blackish	Yellowish
2	Odour	Foul-pungent	Stinking-pungent
3	Transparency	Blackish	Yellowish
4	Turbidity (NTU)	14.0 ± 0.0	8.5 ± 0.07
5	Temperature ($^\circ\text{C}$)	26.66 ± 1.0	32.33 ± 1.54
6	pH	6.66 ± 0.098	7.5 ± 0.1
7	Conductivity (dsm^{-1})	630.0 ± 2.46	840.0 ± 0.12
8	Free CO_2 (mg/l)	170.0 ± 0.0	220.00 ± 1.12
9	DO (mg/l)	45.0 ± 1.0	76.33 ± 0.78
10	BOD (mg/l)	188.33 ± 0.78	200.0 ± 1.75
11	COD (mg/l)	370.0 ± 1.0	396.33 ± 0.66
12	TSS (mg/l)	1032.66 ± 2.33	1312.0 ± 2.18
13	TDS (mg/l)	1902.33 ± 2.4	2071.33 ± 2.4
14	Cl- (mg/l)	620.66 ± 1.0	660.33 ± 1.75
15	Ca^{2+} (mg/l)	240.33 ± 1.66	275.33 ± 3.09
16	Na^+ (mg/l)	260.2 ± 5.3	238.66 ± 1.06

17	K ⁺ (mg/l)	53.0 ± 1.05	82.0 ± 1.35
18	Mg ²⁺ (mg/l)	213.33 ± 1.76	265.0 ± 2.51
19	SO ₄ ⁻⁻ (mg/l)	303.0 ± 1.6	403.66 ± 1.56
20	Total alkalinity (mg/l)	170.0 ± 1.52	307.33 ± 2.72
21	Inorganic P (mg/l)	0.16 ± 0.0	0.18 ± 0.0
22	Particulate P (mg/l)	0.52 ± 0.08	1.0 ± 0.0
23	Total Dissolved P (mg/l)	0.76 ± 0.01	1.0 ± 0.0
24	Total Kjeldhal P (mg/l)	2.41 ± 0.01	1.65 ± 0.0
25	NH ₃ -N (mg/l)	1.02 ± 0.01	1.1 ± 0.02
26	NO ₃ -N (mg/l)	18.0 ± 1.45	289.66 ± 1.2

Table 1: Physico-Chemical Characterization of Sewage Effluents from Selected Sites in Mathura.

Values are arithmetic mean ± SEM of three replicates*, **Significant at 0.05% P and 0.01% P respectively; ns- non significant

- Colour:** The water sample showed a Black colour at Site I while Yellowish colour at Site II.
- Odour:** It was found to be Foul-pungent smell at Site I and Stinking-pungent smell at Site II.
- Transparency:** Water transparency also differed at the two sites with Blackish at Site I and 4. Yellowish at Site II.
- Turbidity:** Coming to water turbidity with higher value at Site I (14.0 ± 0.0 NTU) and lower at Site II (8.5 ± 0.07 NTU).
- Temperature:** Site I (26.66 ± 1.0°C) and Site II (32.33 ± 1.54°C).
- Hydrogen Ion Concentration:** Hydrogen ion concentration i.e., pH showed lower values at Site I (acidic 6.66 ± 0.09) whereas at Site II the pH was observed to be (alkaline 7.5 ± 0.1).
- Conductivity:** Conductivity was found to be very high with lower (630.0 ± 2.46 dsm⁻¹) at Site I and higher (840.0 ± 0.12 dsm⁻¹) at Site II.
- Free Carbon Dioxide:** Free Carbon dioxide showed the lower value at Site I (170.0 ± 0.0 mg/l) and highest values were recorded at Site II (220.0 ± 1.12 mg/l).
- Dissolved Oxygen:** Observations of Dissolved oxygen (DO) levels showed lower values at Site I (45.0 ± 1.0 mg/l) with an upward trend at Sites II (76.33 ± 0.78 mg/l).
- Biological Oxygen Demand:** Biological Oxygen Demand (BOD) levels also showed very high values with lower at Site I (188.33 ± 0.78 mg/l) with an upward trend at Site II (200.0 ± 1.75 mg/l).
- Chemical Oxygen Demand:** Chemical Oxygen Demand (COD) values were also observed to be high enough ranging from lower values at Site I (370.0 ± 1.0 mg/l) followed with upward values at Sites II (396.33 ± 0.66 mg/l).
- Total Suspended Solids:** Total Suspended Solids (TSS) with lower values at Site I (1032.66 ± 2.33 mg/l) and the higher at the Site II (1312.0 ± 2.08 mg/l).
- Total Dissolved Solids:** Total Dissolved Solids (TDS) content was lower at Site I (1902.33 ± 2.4 mg/l) and higher at Site II (2071.33 ± 2.4 mg/l).
- Chloride Ions:** Regarding the Anion (-ve ions) Chloride ions (Cl⁻) the variation was quite observable with lower being at Site I (620.66 ± 1.0 mg/l) followed upwardly at the Sites II (660.33 ± 1.75 mg/l).
- Calcium Ions:** Calcium ions (Ca²⁺) at Site I (240.33 ± 1.66 mg/l) were lower followed upwardly at the Site II (275.33 ± 3.09 mg/l).
- Sodium Ions:** Sodium ions (Na⁺) at Site I (260.2 ± 5.3 mg/l) were the higher than at the Site II (238.66 ± 1.06 mg/l).
- Potassium Ions:** Potassium ions (K⁺) at Site I (53.0 ± 1.05 mg/l) followed upwardly at the Site II (82.0 ± 1.35 mg/l).
- Magnesium Ions:** Magnesium ions (Mg²⁺) at Site I (213.33 ± 1.76 mg/l) followed upwardly at the Site II (265.0 ± 2.51 mg/l).
- Sulphate Ions:** Sulphate ions (SO₄²⁺) at Site I (303.0 ± 1.6 mg/l) followed upwardly at the Site II (403.66 ± 1.56 mg/l).
- Total alkalinity:** Total alkalinity at two sites was found to be at Site I (179.0 ± 1.52 mg/l) and at Site II (315.33 ± 2.72 mg/l).

21. **Inorganic Phosphate:** Inorganic phosphate at two sites was found to be at Site I (0.17 ± 0.0 mg/l) and at Site II (0.19 ± 0.0 mg/l).
22. **Particulate Phosphate:** Particulate P at two sites was found to be at Site I (0.53 ± 0.08 mg/l) and at Site II (1.2 ± 0.0 mg/l).
23. **Total Dissolved Phosphate:** Total Dissolved P at two sites was found to be at Site I (0.81 ± 0.01 mg/l) and at Site II (1.1 ± 0.0 mg/l).
24. **Total Kjeldhal Phosphate:** Total Kjeldhal P at two sites was found to be at Site I (2.62 ± 0.01 mg/l) and at Site II (1.6 ± 0.0 mg/l).
25. **Ammonium Nitrate:** $\text{NH}_3\text{-N}$ at two sites was found to be at Site I (1.12 ± 0.01 mg/l) and at Site II (1.24 ± 0.02 mg/l).
26. **Nitrate Nitrogen:** $\text{NO}_3\text{-N}$ at two sites was found to be at Site I (20.33 ± 1.45 mg/l) and at Site II (311.66 ± 1.2).
27. Statistically the two sites showed highly significant differences at both 0.01% and 0.05% level of probability conforming to high pollution at the two sites.

The sources of odourous substances in potable water have been recorded by Sayato [1976], Singh [2003], Sharma [2004], Singh., *et al.* [2008], Parashar [2011]; Sharma., *et al.* [2013a, b]; Snehlata [2016]; Parashar and Sharma [2017]; Parashar., *et al.* [2018]; Sharma., *et al.* [2018]. As such water quality in River Yamuna observed showed foul-stinking to pungent smell owing to higher levels of TDS, chlorides, sulphates, SS and total hardness. In the present investigation it was found that most of the industries (printing, dyeing, silver polishing, supari, chemical industries etc.) discharge their effluents in the Yamuna River through various drains, therefore, the two sites studied showed different colours in the water. The objectionable colour observed at the two sites was mainly due to the addition of city sewage as well as industrial effluents discharged at these sites.

The description presented in the present investigation reveals that the transparency varied with from bluish-black to yellowish-green tinge as was also revealed in the earlier studies [Thakur, 2001; Sharma, 2004; Parashar, 2011; Sharma., *et al.* [2013a, b]; Snehlata, 2016; Parashar., *et al.* 2018; Sharma., *et al.* [2018].

The suspended solids increase the turbidity of river water. The maximum permitted turbidity limit by set by WHO [1984] for drinking water is 5 – 8 NTU and for other domestic purposes that should not exceed more than 25 NTU. Turbidity showed positive relation with temperature, BOD, COD, DS and SS. However, the negative correlation was found with DO and the nitrates as has also been reported by a number of workers [Singh., *et al.* 1989a, b; Murthy., *et al.* 1994; Thakur, 2001; Singh, 2003; Sharma, 2004; Singh., *et al.* 2008; Parashar, 2011; Sharma., *et al.* [2013a, b]; Snehlata, 2016; Parashar., *et al.* 2018; Sharma., *et al.* [2018]. The present study is in conformity of with the earlier findings [Rai, 1974a, b; Mitra, 1982; Yadav and Saxena, 1987; Sehgal and Siddiqui, 1989; Thakur, 2001; Sharma, 2004; Parashar, 2011; Sharma., *et al.* [2013a, b]; Snehlata, 2016; Parashar., *et al.* 2017; Sharma., *et al.* [2018] which reported water temperature variations contaminated by effluents from various types of industries. Our results clearly revealed that the presence of sewage effluents from city and industrial discharges is the major cause of rising temperature. With the rise of temperature there is decline in the DO levels. Also a rise in temperature leads to the speeding up of the physico-chemical reactions in water reducing the solubility of gases as such temperature showed strong positive correlation with turbidity, BOD, COD, DO, TSS and TDS and a mild correlation with pH, ammonia and nitrates. The strong correlation of temperature with turbidity, DS and SS is due to the quick evaporation of water when temperature is high. In concentrated water their density becomes high and leads to high turbidity as is shown by a number of workers [Thakur, 2001; Sharma, 2004; Parashar, 2011; Sharma., *et al.* [2013a, b]; Snehlata, 2016; Parashar., *et al.* 2018; Sharma., *et al.* [2018].

The pH is another most important factor that influences the rate of biochemical conversion. Further, pH showed its positive correlation with all the parameters except the DO which might probably be due to that oxygen reduces the alkalinity of ammonical wastes [Dakshini and Soni, 1979; Mathur., *et al.* 1987; Singh., *et al.* 1988; Raiyani., *et al.* 1994; Thakur, 2001; Singh, 2003; Sharma, 2004; Singh., *et al.* 2008; Parashar, 2011; Sharma., *et al.* [2013a, b]; Snehlata, 2016; Parashar., *et al.* 2018; Sharma., *et al.* [2018].

According to WHO [1984] the DO should not be lower than 6 mg/l (3 mg/l ISI standard). During present study the DO showed a highly fluctuating trend as it exhibited a strong negative trend with temperature, BOD, COD, ammonia, DS and SS. Depletion of DO is an index of increased organic pollution. The water quality in respect to DO suggests that both the Sites I and II were highly polluted. The

maximum permissible limit for BOD is 5 mg/l [WHO, 1984]. BOD was found to be very high at Site II. It was found to have a very strong negative correlation with DO. High COD shows high accumulation of organic waste in the river. Broadly COD exhibited similar trends to BOD. COD was high at both the sites. *Free Carbon dioxide showed the higher values*. COD values were also observed to be *high enough*.

The above findings clearly showed undisputed fact that trends in DO, BOD and COD levels remained uniform and did not differ to a larger extent. Self-purification of rivers is one of their most remarkable character which leads eventually to a large extent on biochemical reactions going on in the water brought about by the activities of micro-organisms (bacteria and phytoplanktons) which replenish sufficient oxygen utilizing the organic matter as food and breakdown complex compounds to simpler and harmless products. The determination of BOD at various points in the river is necessary to assess the extent to which self-purification has proceeded.

Total dissolved solids (TDS) also serve as indicator of sewage effluents which adds to turbidity and therefore, positively correlated with turbidity along with temperature, BOD and COD [Saxena and Chauhan, 1993; Thakur, 2001; Singh, 2003; Sharma, 2004; Singh, *et al.* 2008; Parashar, 2011; Sharma, *et al.* [2013a, b]; Snehlata, 2016; Parashar, *et al.* 2018; Sharma, *et al.* [2018]]. TDS also showed positive trend with hardness. Since the ground water of Mathura depicts higher values of hardness, this might be the reason for high TDS. Comparatively TDS was higher due to the continuous mixing of industrial effluents. Total suspended solids (TSS) showed almost a similar trend as that of TDS and were also found to be positively correlated with turbidity along with temperature, BOD and COD and hardness but showed a negative relation with that of DO. There seems to be an indirect negative relationship between TSS and DO involving the role of temperature and turbidity [Datar and Vashistha, 1992; Sunderrajan, *et al.* 1994; Thakur, 2001; Singh, 2003; Sharma, 2004; Singh, *et al.* 2008; Parashar, 2011; Sharma, *et al.* 2013a, b; Snehlata, 2016; Parashar, *et al.* 2017; Sharma, *et al.* 2018]. TSS was, however, higher at Sites I and II probably due to the accumulation of large amount of un-decomposed organic and inorganic wastes indicating positive correlation of TSS with BOD which is in corroboration with the observations made by Mathur, *et al.* [1987], Saxena and Chauhan, [1993], Shaji and Patel, [1990], Thakur [2001], Singh [2003], Sharma [2004], Singh, *et al.* [2008], Parashar [2011], Sharma, *et al.* [2013a, b]; Snehlata [2016], Parashar, *et al.* [2017]; Sharma, *et al.* [2018]. Further, analysis of water revealed the presence of maximum amounts of electrolytes as such the conductivity was found to be very high. Regarding the Anion (-ve ions) Chloride ions (Cl⁻) the variation was quite observable at the two sites whereas a reverse trend was observed with cations, i.e., Ca²⁺, Mg²⁺, K⁺ and Na⁺, where the maximum amount was found in water samples of Site II. The other ions like SO₄²⁻ matched with concentration of cations. Likewise, Cations (+ve ions) Calcium ions (Ca²⁺), Sodium ions (Na⁺), Potassium ions (K⁺) and Magnesium ions (Mg²⁺) also showed variable trends. The probable reason for the high degree of salinity might be due to the existing higher levels of salts of Ca, Mg, Na and K in the ground water along with the various chemical industries discharging their effluents and huge quantity of untreated city sewage in river water [Saxena and Chauhan, 1993; Thakur, 2001; Singh, 2003; Sharma, 2004; Singh, *et al.* 2008; Gautam, 2010; Parashar, 2011; Snehlata, 2016; Parashar, *et al.* 2017]. The total organic and inorganic solids present in industrial and sewage effluents affect the turbidity, conductivity and alkalinity, as highest amount of TS was observed a greater quantum of total solids has shown resemblance with above discussed parameters. As recorded earlier a similar finding was observed with TDS where a higher magnitude of TDS was recorded at Site II surprisingly enough followed by Site I because a huge amount of effluents are directly thrown by various industries and city sewage through a number of drains.

Seed Germination

ANOVA Analysis

The results in relation to the effect of different concentrations of Sewage Effluent (Sewage Farm) exposure on germination performance measured in terms of Germination Percent (GP) after 24, 48, 72, 96 and 120 hours of sowing have been shown in TABLES 2 – 11. As indicated in the Table 2 (ANOVA ANALYSIS - Germination Percent- GP) all the main effects viz., Crop, Treatment, Duration, and their interactions (A X B; A X C; B X C; A X B X C; A X D; B X D; A X B X D; C X D; A X C X D; B X C X D and A X B X C X D) were found to be highly significant at 0.01 and 0.05% level of probability with significant differences noticed in the GP of all the crops studied TABLES 2 – 11.

Source of variation	DF	SS	MSS	F-value	Significance
Factor A (Crop)	3	4055.99	1351.99	631.88	*,**
Factor B (Treatment)	4	73941.92	18485.48	8639.6	*,**
Factor C (Duration)	4	151578.4	37894.61	17710.9	*,**
Factor A X B	12	18077.8	1506.48	704.08	*,**
Factor A X C	12	43626.52	3635.54	1699.15	*,**
Factor B X C	16	13995.17	874.69	408.81	*,**
Factor A X B X C	48	8792.95	183.18	85.61	*,**
Error	200	427.92	2.14		
Total	299	496.7			
SEM ± 0.08 Grand Mean = 42.96					
Range of Germination	Lower Range = 42.8 Upper Range = 43.13				

Table 2: *Anova Table *Effect of Different Levels of Sewage Effluent Exposure on Germination Percentage in the Four Crops.

***Significant at 0.05% Level of Probability; **Significant at 0.01% Level of Probability**

Interaction Crop: The highest mean GP (47.04%) was recorded in the crop C3 followed by C1, C2 and lastly C4 with the lowest germination percentage of (39.29%). The crops were arranged in the following descending order on the basis of their respective GP irrespective of the treatment and seedling age (interaction Crop) (Table 3): C3 Mustard (47.04%) > C1 Wheat (46.22%) > C2 Barley (39.31%) > C4 Broad bean (39.29%)

Crop	Germination Percentage	Range of Germination Percentage	
		Lower	Upper
C1 Wheat	46.22	45.89	46.56
C2 Barley	39.31	38.98	39.64
C3 Mustard	47.04	46.70	47.37
C4 Broad Bean	39.29	38.96	39.62
SEM ± 0.169			

Table 3: Effect of Different Levels of Sewage Effluent Exposure on Germination Percentage in the Four Crops (Crop).

Interaction Treatment: A significant reduction in GP with increasing SE levels was observed irrespective of variety, treatment and seedling age (interaction Treatment). *The reduction in GP was more pronounced after 75% of SE treatment.* Further, with treatment GP was highest in controls (61.105%) followed by 50% SE (52.01%) showing increase over 25% SE while lowest (17.001%) with 100% SE (TABLE 4) and the treatments were arranged in the following descending order: Control (61.10%) > 50% (52.01%) > 25% (50.92%) > 75% (33.8%) > 100% (17.0%)

Treatment Sewage Effluent (%)	Germination Percentage	Range of Germination Percentage	
		Lower	Upper
Control DW	61.1	60.73	61.47
25%	50.92	50.55	51.29
50%	52.01	51.64	52.38
75%	33.8	33.43	34.17
100%	17.0	16.62	17.37
SEM ± 0.189			

Table 4: Effect of Different Levels of Sewage Effluent Exposure on Germination Percentage in the Four Crops (Treatment).

Interaction Duration: It was observed that the crops showed first sign of germination at 24 hrs after sowing irrespective of crops and SE treatment (interaction Duration) and thereafter GP increased significantly with seedling age till 120 hrs (TABLE 5) and the treatments were arranged in the following descending order: 120 hr (66.32%) > 96 hr (60.87%) > 72 hr (53.95%) > 48 hr (26.19%) > 24 hr (7.49%)

Duration (hours)	Germination Percentage	Range of Germination Percentage	
		Lower	Upper
24 hrs	7.49	7.12	7.87
48 hrs	26.19	25.82	26.56
72 hrs	53.95	53.58	54.32
96 hrs	60.87	60.50	61.25
120 hrs	66.32	65.94	66.69
SEM ± 0.189			

Table 5: Effect of Different Levels of Sewage Effluent Exposure on Germination Percentage in the Four Crops (Duration).

Interaction Crop X Treatment: The significant interaction of varieties with treatment (Crop X Treatment) is depicted in TABLE 6. All the four varieties showed a decrease in GP with increasing SE treatment however, the genotypic variations were quite evident. Surprisingly in 50% SE in leguminous crops C3 mustard and C4 broad bean enhancement in GP over Control was observed. Also in cereal crop C1 wheat in 50% SE treatment enhancement in GP over 25% SE was reported. Further, highest and lowest GP was observed in cereal crop C2 Barley as in Controls (80.318%) and 100% SE (1.275%). The four crops were arranged in the following descending order on the basis of their respective GP irrespective of the treatment:

C1 Wheat – Control (59.72%) > 50% (58.66) > 25% (50.83%) > 75% (36.39%) > 100% (25.52%)
 C2 Barley – Control (80.31%) > 25% (47.95) > 50% (40.98%) > 75% (26.03%) > 100% (1.27%)
 C3 Mustard – 50% (56.19) > 25% (52.9%) > Control (52.37%) > 75% (41.51%) > 100% (32.2%)
 C4 Broad Bean – 50% (52.2) > Control (52%) > 25% (52%) > 75% (31.26%) > 100% (9%)
 C3 Mustard > C1 Wheat > C4 Broad bean > C2 Barley

Treatment Sewage Effluent (%)	Crops			
	C1 Wheat	C2 Barley	C3 Mustard	C4 Broad Bean
Control DW	59.72	80.31	52.37	52.0
25%	50.83	47.95	52.90	52.0
50%	58.66	40.98	56.19	52.2
75%	36.39	26.03	41.51	31.26
100%	25.52	1.27	32.20	9.0
SEM ± 0.378				

Table 6: Effect of Different Levels of Sewage Effluent Exposure on Germination Percentage in the Four Crops (Crop X Treatment).

Interaction Crop X Duration: All the crop varieties showed an increase in GP (interaction Crop X Duration) exhibiting marked differences in their early seedling growth with advancement in seedling age and the effect of SE treatment, i.e., in general, tolerance to SE increased (Table 7) and were arranged in the following descending order on the basis of their respective GP irrespective of the duration:

C1 Wheat – 120 hr (76.66%) > 96 hr (67.98%) > 72 hr (53.57%) > 48 hr (32.66%) > 24 hr (0.26%)
 C2 Barley – 120 hr (42.56%) > 72 hr (42.46%) > 96 hr (42.4%) > 48 hr (39.44%) > 24 hr (29.68%)
 C3 Mustard – 120 hr (79.98%) > 96 hr (67.92%) > 72 hr (54.57%) > 48 hr (32.66%) > 24 hr (0.0%)
 C4 Broad Bean– 120 hr (66.06%) > 96 hr (65.2%) > 72 hr (65.2%) > 48 hr (0.0%) > 24 hr (0.0%)
 C3 Mustard > C1 Wheat > C4 Broad bean > C2 Barley

Duration (hours)	Crops			
	C1 Wheat	C2 Barley	C3 Mustard	C4 Broad Bean
24 hrs	0.26	29.68	0.0	0.0
48 hrs	32.66	39.44	32.66	0.0
72 hrs	53.57	42.46	54.57	65.2
96 hrs	67.98	42.4	67.92	65.2
120 hrs	76.66	42.56	79.98	66.06
SEM ± 0.378				

Table 7: Effect of Different Levels of Sewage Effluent Exposure on Germination Percentage in the Four Crops (Crop X Duration).

Interaction Duration X Treatment: The interaction of Duration X Treatment (Table 8) shows that with increasing SE treatment level the deleterious salt effect was clearly observable which, however, declined with seedling age. Initially at 24 hours seedling age in control sets showed 12.667% germination but no germination was observed in the SE treatment sets of 100%. The treatments were arranged in the following descending order:

Control - 120 hr (89.06) > 96 hr (85.12) > 72 hr (76.75) > 48 hr (41.92) > 24 hr (12.66)
 25% - 120 hr (77.58) > 96 hr (72.5) > 72 hr (64.38) > 48 hr (30.02) > 24 hr (10.12)
 50% - 120 hr (78.59) > 96 hr (73.69) > 72 hr (67.1) > 48 hr (19.85) > 24 hr (8.96)
 75% - 120 hr (55.91) > 96 hr (47.15) > 72 hr (40.37) > 48 hr (35.75) > 24 hr (5.71)
 100% -120 hr (30.45) > 96 hr (25.91) > 72 hr (21.15) > 48 hr (7.45) > 24 hr (0.0)

Treatment Sewage Effluent (%)	Duration (hours)				
	24 hrs	48 hrs	72 hrs	96 hrs	120 hrs
Control DW	12.66	41.92	76.75	85.12	89.06
25%	10.12	30.02	64.38	72.50	77.58
50%	8.96	31.70	67.10	73.69	78.59
75%	5.71	19.85	40.37	47.15	55.91
100%	0.00	7.45	21.15	25.91	30.45
SEM ± 0.422					

Table 8: Effect of Different Levels of Sewage Effluent Exposure on Germination Percentage in the Four Crops (Duration X Treatment).

Interaction Crop X Duration X Treatment: A review of the final interaction (Crop X Duration X Treatment) reveals that irrespective of salt concentration levels the GP had increased with seedling age and that the SE treatment had their individual effect depending upon the varying treatment levels. The GP was significantly inhibited by SE treatment in all the four crops however the degree of inhibition varied depending on the concentration of SE treatment (Tables 1–10). The lowest and highest at each duration from 24 to 120 hours and each treatment have been marked with lowest (*) and highest (**) asterisk in the Table 9 with lowest as 0.0% and highest 95.23 at 120 hours in 50% SE treatment in the crop C1 wheat. The lowest (0.0%) and the highest (57.67%) germination in 100% SE treatment have been reported in C3, C4 (lowest) and C3 (highest) respectively. At some hours in crops C1, C3 and C4 enhancement over the control have been observed in 25% and 50% SE treatment as in C3 at 48 hours in 25% SE treatment; in C1 at 24, 72, 96 and 120 hours in 50%; in C3 at 48 to 120 hours in 50% and in C4 from 72 to 120 hours in 50% SE treatment.

Crops	Duration (hours)	Treatment Sewage Effluent (%)				
		Control DW	25%	50%	75%	100%
C1 Wheat	24 hrs	0.37	0.35	0.43	0.32	0.05
	48 hrs	52.02	35.28	38.88	25.50	11.62
	72 hrs	67.95	58.96	71.90	38.45	30.60
	96 hrs	86.95	75.30	86.90	52.45	38.34
	120 hrs	91.35**	84.30	95.23**	53.79	47.01
C2 Barley	24 hrs	50.02**	40.15**	35.43**	22.75**	0.08**
	48 hrs	82.40**	46.82**	41.34	26.11	0.53
	72 hrs	89.70**	51.05*	42.70*	27.24*	1.67*
	96 hrs	89.75**	50.56*	42.72*	27.24*	1.98*
	120 hrs	89.75	51.19*	42.75*	27.71*	2.12*
C3 Mustard	24 hrs	0.25	0.00*	0.00*	0.00*	0.00*
	48 hrs	33.27*	37.99	46.62**	27.81**	17.67**
	72 hrs	62.73*	60.85	66.81	45.15	37.34**
	96 hrs	77.13*	77.52	78.15	58.48**	48.34**
	120 hrs	88.50	88.18**	89.40	76.15**	57.67**

C4 Broad Bean	24 hrs	0.00*	0.00*	0.00*	0.00*	0.00*
	48 hrs	0.00*	0.00*	0.00*	0.00*	0.00**
	72 hrs	86.67	86.67**	87.00**	50.67**	15.00
	96 hrs	86.67	86.67**	87.00**	50.67	15.00
	120 hrs	86.67	86.67	87.00	55.00	15.00
SEM ± 0.845						

Table 9: Effect of Different Levels of Sewage Effluent Exposure on Germination Percentage in the Four Crops (Crop X Duration X Treatment).

*, **show lowest and highest values at each duration respectively

Seeing overall results it was recorded that the lowest GP (0.0%) was found in 25%, 50%, 75% and 100% SE treatment at 24 hours of germination in the crop C3 and C4 at 24 and 48 hours in controls as well as 25%, 50%, 75% and 100% SE treatment whereas highest (95.23%) was recorded in 50% SE treatment at 120 hours in C1. Also it has been observed that the crops C1 and C3 behaved better, even in 100% SE treatment concentration at 120 hours of seedling growth. Thus, crops show overall GP as: C3 Mustard > C1 Wheat > C4 Broad bean > C2 Barley

Germination percentage as percent over control showed lowest rate (0.0%) at 24 hours in 100% in the four crops and the highest of (140.1300%) at 48 hours in 50% in the crop C3 (TABLE 10). On the basis of percent over control varieties were placed as: C3 Mustard > C1 Wheat > C4 Broad bean > C2 Barley

Crops	Duration (hours)	Treatment Sewage Effluent (%)				
		Control DW	25%	50%	75%	100%
C1 Wheat	24 hrs	100.0	94.59	116.21	28.91	13.51
	48 hrs	100.0	67.83	74.75	49.02	22.33
	72 hrs	100.0	86.77	105.81	56.58	45.03
	96 hrs	100.0	86.60	99.94	60.32	44.22
	120 hrs	100.0	92.28	104.24	47.93	51.46
C2 Barley	24 hrs	100.0	80.27	70.82	45.47	0.00
	48 hrs	100.0	56.81	56.16	31.68	0.64
	72 hrs	100.0	56.91	47.60	30.36	1.85
	96 hrs	100.0	56.33	47.60	30.11	2.22
	120 hrs	100.0	57.03	47.63	30.13	2.35
C3 Mustard	24 hrs	100.0	0.00	0.00	0.00	0.00
	48 hrs	100.0	114.18	140.13	83.59	53.10
	72 hrs	100.0	97.00	106.50	71.97	59.52
	96 hrs	100.0	100.49	101.31	75.82	62.67
	120 hrs	100.0	99.64	101.01	86.04	65.16

C4 Broad Bean	24 hrs	100.0	0.0	0.00	0.00	0.00
	48 hrs	100.0	0.0	0.00	0.00	0.00
	72 hrs	100.0	100.0	100.384	58.462	17.308
	96 hrs	100.0	100.0	100.384	58.462	17.308
	120 hrs	100.0	100.0	100.384	63.461	17.308
SEM ± 0.845						

Table 10: Effect of Different Levels of Sewage Effluent Exposure on Germination Percentage in the Four Crops (Percent Over Control) (Crop X Duration X Treatment).

Data recorded after 72 hours of presoaking the seeds in test solution have shown highest rate of GP (percent over control) 106.509 (+ 6.5) in 50% SE treatment in the crop C3 and lowest 1.85 (-98.14) in 100% in C2 depict C3 growth of + 6.5% over the control and the overall better performance of the C3 in all the treatments (TABLE 11) and were arranged in the following descending order:

C1 Wheat – 50% [105.81 (+ 5.81)] > Control [100%] > 25% [86.77 (- 13222)] > 75% [56.58 (- 43.41)] > 100% [45.03 (- 54.96)]
 C2 Barley –Control [100%] > 25% [56.91 (- 43.09)] > 50% [47.6 (- 52.39)] > 75% [30.36 (- 69.63)] > 100% [1.85 (- 98.14)]
 C3 Mustard –50% [106.5 (+ 6.5)] > Control [100%] > 25% [97.0 (- 2.99)] > 75% [71.97 (- 28.02)] > 100% [59.52 (- 40.47)]
 C4 Broad Bean – 50% [100.38 (+ 0.38)] > Control [100%] > 25% [100.0 (- 0.0)] > 75% [58.462 (- 41.538)] > 100% [17.3 (- 82.69)]

Crop	Treatment Sewage Effluent (%)				
	Control DW	25%	50%	75%	100%
C1 Wheat	100	86.77 *(-13.22)	105.81 *(+5.81)	56.58 *(-43.41)	45.03 *(-54.96)
C2 Barley	100	56.91 *(-43.09)	47.6 *(-52.39)	30.36 *(-69.63)	1.85 *(-98.14)
C3 Mustard	100	97.00 *(-2.99)	106.5 *(+6.5)	71.97 *(-28.02)	59.52 *(-40.47)
C4 Broad Bean	100	100.00 *(-0.0)	100.38 *(+0.38)	58.46 *(-41.53)	17.3 *(-82.69)
SEM ± 0.845					

Table 11: Effect of Different Levels of Sewage Effluent Exposure on Germination Percentage in the Four Crops Percent over Control.

(Data Recorded After 72 Hours of Presoaking the Seeds in Test Solutions)

*Values in Parenthesis show Percent Increase (+)/Decrease (-) over Control in GP

Further, it was observed that 75% SE concentrations and also 72 hours of germination were found to be quite critical for the crops, therefore, on the basis of percent over control at 72 hours with 75% SE concentration the germination percentage in the four crops showed a trend as percent change over control: Mustard [71.97%], Broad bean [58.46], Wheat [56.58] and Barley [30.36] showing better performance of the leguminous crops than the cereal crops (Table 11). Finally on the basis of percent germination over control in critical level of 75% at 72 hours the crops were arranged in the following descending order where the two leguminous crops showed better results than the two cereal crops: GP: C3 Mustard [71.97] > C4 Broad bean [58.46] > C1 Wheat [56.58] > C2 Barley [30.36]

Speed of germination index (SGI)

Lastly, the results for Speed of Germination Index (SGI) have been depicted in the TABLE 12. Both the legume crops (C3 Mustard and C4 Broad bean) at 25% and 50% SE treatment concentrations showed an increase (+) in GP of treated seeds over control but at higher concentrations of 75% and 100% SE treatment concentration a decrease (-) was reported in the two crops. In the crop C3

increase was found to be 2.2% in 25% concentration and at 50% concentration it rose to 11.91% whereas in crop C4 at 25% concentration it was same as that of control and at 50% concentration GP was slightly higher (0.38% over control). At higher concentrations of 75% and 100% SE treatment in the legume crop C3 decrease (-) of GP was found to be 22.19% and 40.37% over control respectively. In the other legume crop C4 decrease (-) of GP at higher concentrations of 75% and 100% SE treatment was found to be quite higher in that 40.7% and 82.693% over control respectively (Table 12). On the other hand, in the two cereal crops (C1 wheat & C2 Barley) there was no increase (+) in GP was observed and also the crops showed absolutely differential pattern in that in crop C2 decrease (-) was found to be quite sharp as compared to the crop C1. Thus, overall SGI results showed highest (111.91% over control) in the legume crop C4 Mustard at 50% SE treatment concentration whereas the lowest (1.22% over control) SGI was observed in the cereal crop C2 Barley at 100% SE treatment concentration (Table 12).

Crop	Speed of germination index (SGI) Percent increase (+)/decrease (-) over Control				
	Treatment Sewage Effluent (%)				
	Control DW	25%	50%	75%	100%
C1 Wheat	100	81.68 *(-18.31)	94.6 *(-5.39)	55.61 *(-44.38)	38.61 *(-61.38)
C2 Barley	100	62.02 *(-37.97)	53.55 *(-46.44)	34.17 *(-65.82)	1.22 *(-98.77)
C3 Mustard	100	102.2 *(+2.2)	111.91 (+11.91)	77.8 *(-22.19)	59.62 *(-40.37)
C4 Broad Bean	100	100.0 *(± 0.0)	100.38 *(+0.38)	*59.29 (-40.7)	17.3 *(-82.69)

Table 12: Effect of Different Levels of Sewage Effluent Exposure on Speed of Germination Index (Sgi) In the Four Crops Percent over Control.

*Values in Parenthesis show Percent Increase (+)/Decrease (-) over Control in GP

Further, it was observed that 75% SE concentrations and also 72 hours of germination were found to be quite critical for the crops, therefore, on the basis of percent over control at 72 hours with 75% SE concentration the germination percentage in the four crops showed a trend as percent change over control: Mustard [71.8%], Broad bean [59.29], Wheat [55.61] and Barley [34.17] showing better performance of the leguminous crops than the cereal crops. Finally, on the basis of SGI over control in critical level of 75% at 72 hours the crops were arranged in the following descending order where the two leguminous crops showed better results than the two cereal crops: SGI: C3 Mustard [71.8] > C4 Broad bean [59.29] > C1 Wheat > [55.61] > C2 Barley [34.17]

Thus, overall results of GP and SGI proved that the two legume crops Mustard and Broad bean showed more tolerant behavior than the cereal crops Wheat and Barley towards Sewage Effluent irrigation with dilution in the concentration as 25%, 50% and 75% and that there was found to be a direct correlation between GP and SGI at critical SE level of 75% and duration 72 hours as follows (Table 13):

Crops	Germination Percentage (GP)	Speed of Germination Index (SGI)
C3 Mustard	71.97	77.8
C4 Broad bean	58.46	59.29
C1 Wheat	56.58	55.61
C2 Barley	30.36	34.17

Table 13: Effect of Different Levels of Sewage Effluent Exposure on Germination Percentage (Gp) and Speed of Germination Index (Sgi) In the Four Crops Percent Increase (+)/Decrease (-) Over Control at 72 Hours and in 75% Sewage Effluent (Se) Concentration.

GP: C3 Mustard [71.97] > C4 Broad bean [58.46] > C1 Wheat [56.58] > C2 Barley [30.36]

SGI: C3 Mustard [71.8] > C4 Broad bean [59.29] > C1 Wheat [55.61] > C2 Barley [34.17]

Increasing urbanization and industrialization is posing the disposal of sewage effluents on agricultural land a serious problem all over the world [Parashar and Sharma, 2017]. The continuous use of industrial effluents by people in the adjoining area has become a common practice which results in accumulation of industrial pollutants and toxic metals in soil to such a level that may become phytotoxic and hazardous for the life of animal and human beings [Singh, 2009]. The polluted sewage effluent water inhibits enzyme action thus affecting first of all the first step of plant life i.e., germination of seeds [Shani, *et al.* 1983; Parashar, 2011; Sharma, *et al.* 2014a, b, c, 2016; Sharma, 2015, 2017; Snehlata, 2016; Parashar and Sharma, 2017]. Besides, polluted sewage effluents also contain beneficial elements (N, P, K) as well as toxic metals (Cr, Ni, Zn, Cd, Pb, Hg, Pb and As) as reported earlier by several researchers [Mishra, *et al.* 1998; Sharma, 2004; Parashar, 2011; Sharma, 2015, 2017; Snehlata, 2016; Parashar, *et al.* 2017]. Also increase in $\text{NH}_4^+\text{-N}$ concentration in effluent due to the mineralization of easily decomposable nitrogenous substances present in the sludge resulting in the release of $\text{NH}_4^+\text{-N}$ and later the decrease in the concentration of $\text{NH}_4^+\text{-N}$ may be due to plant uptake and reduced rate of release of $\text{NH}_4^+\text{-N}$ from sludge [Singh, *et al.* 1992], fixation and volatilization losses [Nagarajah, 1988].

The effluents along with N, P, K released higher amounts of $\text{NH}_4^+\text{-N}$. Besides, $\text{NO}_3^-\text{-N}$ recorded due to withholding of water and conversion of $\text{NH}_4^+\text{-N}$ to $\text{NO}_3^-\text{-N}$ (mineralization). The $\text{NO}_3^-\text{-N}$ concentration may be due to immobilization and denitrification by general soil microflora. A higher $\text{NO}_3^-\text{-N}$ concentration recorded may also be caused by oxidation of $\text{NH}_4^+\text{-N}$ to $\text{NO}_3^-\text{-N}$ in the rhizosphere, resulting due to the excess proliferation of plant roots rendering the soil more oxidative [Sakai, 1970; Smith and Delaune, 1984]. The present study revealed that the higher N content in the effluents released significantly higher N fraction through decomposition which can be considered as an organic manure and soil amendment if properly treated and diluted before being used for irrigating crops. Similarly, the addition of sludge to agricultural land has been reported to increase the yields of many plants [Sharma, 2004; Unnamalai, *et al.* 2007; Parashar, 2011; Sharma, 2015, 2017; Parashar and Sharma, 2017].

In the present analysis investigation was undertaken to assess the effect of sewage effluent on water quality of Yamuna River at Mathura and its impact on germination and later seedling growth and metabolism (physiological as well as well biochemical) of the crop plants such as Wheat, Barley, Mustard and Broad bean crops usually cultivated in this area. In our laboratory several researchers working on different cereal and legume crops have shown similar work for quite a long time with different salts, heavy metals, city and industrial effluents [Sharma, 1982, 1987, 2015, 2017; Sharma, 2004; Nauhbar, 2005; Yadav, 2006; Rani, 2007; Gautam, 2009; Singh, 2009; Gautam, 2010; Parashar, 2011; Snehlata, 2016; Sharma, 2015, 2017; Pathak, 2017; Parashar and Sharma, 2017; Saraswat, 2017]. The effects of city and industrial effluents have been well documented in various crops [Sahai, *et al.* 1983; Rajendran, 1990; Sharma, *et al.* 2000; Sharma, 2004; Indira and Mohanty, 2006; Mahapatro and Mohanty, 2007; Dalai and Mohanty, 2008] suggesting various degrees of dilution of effluents beneficial to the crops. Corroborating the present investigation, studies conducted by various other researchers [Kadioglu and Algur, 1990; Tiwari, *et al.* 1993; Muthukumar and Arokiasamy, 1994; Subramani, *et al.* 1995; Sharma, 2004; Dalai and Mohanty, 2008; Sharma, 2015, 2017; Parashar and Sharma, 2017] revealed that the lower effluent concentration stimulates the plant growth whereas inhibiting at higher concentrations. A large number of reviews [Kannabiran and Pragasam, 1993; Kumar, 1995; Chidambaram, *et al.* 1996; Kannan, 2001; Pragasam and Kannabiran, 2001a, b, 2004; Mishra and Pandey, 2002] are available with regards to the effect of sewage effluents in the laboratory experimentation with various commonly cultivated crop plants. The effects of industrial effluent on germination of seeds were attributed to the presence of bio-toxic substances which alters the seed-water interaction necessary for triggering enzyme activity [Baruah and Das, 1998; Sharma, 2004; Parashar, 2011]. In our laboratory also while working on various crop plants effects of different industrial effluents were also reported by Sharma, *et al.* [2014a] textile waste water on seed germination and growth development of *Vicia faba* L; Sharma, *et al.* [2014b] paper mill effluent on seed germination and seedling growth of *Cyamopsis tetragonoloba* L; Sharma, *et al.* [2014c] paper mill effluent on seed germination and seedling growth of *Vigna radiata* L; Sharma, *et al.* [2016] Paper Mill Effluent on Seed Germination and Seedling Growth of *Vigna radiata* L which

also corroborates present study. The view of retardation in seedling growth due to high concentration of the effluent was also observed by Sahai, *et al.* [1983] in rice, Srivastava [1986] in *Cajanus cajan*, Rajendran [1990] in *Helianthus annuus*, Sharma [2004] in wheat and sorghum. The bio-toxic effect of the effluents on seedlings was evident from the fact that there is overall decrease in germination percentage. The lower concentration of effluent (25%–50%) has a stimulating effect on the seed germination of the crops studied.

The results in relation to the effect of different concentrations of sewage effluent exposure on level of sewage effluent exposure phyto-toxicity measured in terms of Germination Percentage and Seed Germination Index after 24, 48, 72, 96 and 120 hours of sowing have shown the level of sewage effluent exposure phyto-toxicity was significantly increased from control to 25%, 50%, 75% and 100% sewage effluent exposure treatments. The degree of level of sewage effluent exposure phyto-toxicity varied depending on the percentage concentration. Therefore, it is extremely necessary to treat the effluent properly before release to a river system for irrigation in agriculture to minimize harmful impact [Kumar, 1995; Mishra and Pandey, 2002; Sharma, *et al.* 2002; Sharma, 2004; Parashar, 2011; Snehlata, 2016; Parashar, *et al.* 2017]. Studies undertaken by several workers on green gram, black gram, wheat, ragi, mustard, barley, broad bean and sorghum seedlings [Sharma, 2004; Indira and Mohanty; 2006; Mahapatro and Mohanty, 2007; Parashar, 2011; Snehlata, 2016; Parashar and Sharma, 2017] indicated the injury to the crop seedlings at higher effluent concentrations. Overall, it has been observed that the cereal crops suffered more than legume crops.

Concluding Remarks

The aim of this research is to investigate the effects of Sewage Effluents (SE) stress on seed germination, seedling growth and metabolism in the four important crops Wheat, Barley, Mustard and Broad bean commonly cultivated in this region. The effects of SE stress were tested at germination and seedling growth of these crops as these are the key events for the establishment of plants under any prevailing environmental conditions. In general, the germination percentage with increasing concentration levels of the SE treatment was inhibited however, interestingly, stimulation has been observed in growth of the crops as increased germination percentage at moderate levels of Effluent concentration. This stimulation in growth may be attributed to the nutritional supplementation at lower concentration of salts present in the Effluents. However, significant differences were noticed in the germination percentage of the four crops studied with the legume crops Mustard and Broad bean showing better growth performance than the cereal crops Wheat and Barley. Thus, the effluent treatment from early imbibitional period onwards, caused more inhibition in the cereal crops compared to legume crops might be due to induced oxidative damage in membrane which increases the activity of anti-oxidative enzymes. The higher level of effluents caused inhibition of seed germination and activity of anti-oxidative enzymes decreased. Therefore, legume crops were found to be more tolerant than cereal crops. The results in relation to the effect of different concentrations of sewage effluent exposure on germination performance measured in terms of germination percentage (GP) and Speed of Germination Index (SGI) after 24, 48, 72, 96 and 120 hours of sowing have shown significant differences in the germination percentage of all the crops studied with their interactions. Further, data recorded after 72 hours of presoaking the seeds in test solution have shown highest rate of germination (percent over control) 106.5 (+ 6.5) in 50% sewage effluent treatment in the crop C3 and lowest 1.85 (- 98.14) in 100% in C2 depict the overall better performance of the crop C3 in all the treatments. Still, overall results proved that the two legume crops Mustard and Broad bean showed more tolerant behavior than the cereal crops Wheat and Barley towards Sewage Effluent irrigation with dilution in the concentration as 25%, 50% and 75%. Thus, identification of plants which can resist and clean-up metals from soils is required to improve phyto-remediation technologies. Moreover, plants have been proposed as bio-monitors of soil contamination but also as bio-markers of the environmental risk of pollutants [Parashar and Sharma, 2017].

Acknowledgements

The authors wish to thank Principal and Head Departments of Botany and Chemistry K.R. College, Mathura for kindly permitting and helping to work for the present investigation.

References

1. Allen SE., et al. "Chemical Analysis of Ecological Materials". Blackwell Scientific Publications, Oxford (London) (1974):
2. Annan Kofi A. "Water for Life; United Nations International Decade for Action: 2005 – 2015".
3. AOAC. "Official Methods of Analysis of AOAC International 19th Ed". AOAC International, Gaithersburg, Maryland, USA (2012):
4. APHA, AWWA and WPCF. "Standard Methods for Examination of Water and Waste Water". American Public Health Association, American Water Works Association and Water Pollution Control Federation (1995):
5. Astera M. "The Ideal Soil: A Handbook for the New Agriculture". (2014):
6. Baruah BK and M Das. "Study on impact of paper mill effluent on germination behaviour and seedling growth of crop plant *Oryza sativa*". *Pollution Research Paper* 17.1 (1998): 65-68.
7. Burns JC., et al. "Swine lagoon effluents applied to coastal Bermuda grass: 1. Forage yield, quality and element removal". *Journal of Environmental Quality* 14 (1985): 9-14.
8. Campbell CA and HR Davidson. "Effect of temperature, nitrogen fertilization and moisture stress use by Manitou spring wheat". *Canadian Journal of Plant Science* 59 (1983): 603-626.
9. Carley HE and RD Watson. "Effect of various aqueous extract upon seed germination". *International Journal of Plant Sciences* 129 (1968): 57-62.
10. Chaw R and Reves AS. "Effect of waste water on *Mentha piperita* and Spinaceal oleraceae". *Journal of Environmental Biology* 51 (2001): 131-145.
11. Cheraghi M., et al. "Effect of waste water on heavy metal accumulation in Hamedan Province Vegetables". *International Journal of Botany* 5 (2009): 190-193.
12. Chidambaram Pillai S., et al. "Effect of chemical industry waste on germination, growth and some biochemical parameters of *Vigna radiata* L Wilkjeck and *Vigna mungo* L Hepper". *J. Enu. Poll* 3.3 (1996): 131-134.
13. CSE. "Sewage Canal How to Clean the Yamuna; Centre for Science and Environment". New Delhi 2007
14. Dakshini KMM and JK Soni. "Water quality of sewerage drains entering Yamuna at Delhi". *Indian Journal of Environmental Health* 21.4 (1979): 354-360.
15. Dalai Bandana and BK Mohanty. "Ecophysiological studies on sugar mill effluents on the seedling morphology and biochemistry of *Pisum sativum* L". *Plant Archives* 8.1 (2008): 155-158.
16. Datar MO and RP Vashistha. "Physico-chemical aspects of pollution in river Betwa". *Indian Journal of Environmental Protection* 1.8 (1992): 577-580.
17. Ensink HH., et al. "A nation-wide assessment of wastewater use in Pakistan: an obscure activity or a vitally important one?" *Water Policy* 6 (2004): 197-206.
18. Garrard A. "The effect of b-indolyl acetic acid on the germination and root growth of certain members of Cruciferae". *New Phytologist* 53.2 (1945): 165-176.
19. Gautam Aruna. "The problem of saline wastelands and their management – A Biological Approach with special reference to Mathura". Ph. D. Thesis, Dr. B. R. Ambedkar Univ., Agra (2009):
20. Gautam Shelja. "Study of phytotoxicity of heavy metal pollution and its control". Ph. D. Thesis, Dr. B. R. Ambedkar Univ., Agra (2010):
21. Ghimire SK. "Evaluation of industrial effluents toxicity in seed germination and seedling growth of some vegetables". M.Sc. Dissertation, Central Department of Botany (1994), Tribhuvan University (1994):
22. Grob RC. "Chromatographic Analysis of Environment". (1983): 724.
23. Gupta N., et al. "Prevalence of intestinal helminth eggs on vegetables grown in wastewater-irrigated areas of Titagarh, West Bengal, India". *Food Control* 20 (2009): 942-945.
24. Indira Subhadarsini and BK Mohanty. "Effect of Sugar Mill Effluents on Ragi (*Elusine coracona* L) and growth behaviour of its seedlings". *Journal of Animal and Plant Sciences* 19.11 (2006): 463-467.

25. Johnson FLA., *et al.* "Method for Studying Soil-Microflora- Plant Disease Relationship". Burgess Publishing Co., USA (1959):
26. Kadioglu A and OF Algur. "The effect of vinasse on the growth of Helianthus annuus and Pisum sativum Part 1. The effects on some enzymes and chlorophyll and protein contents". *Environmental Pollution* 671 (1990): 223-232.
27. Kannabiran B and A Pragasam. "Effects of distillery effluents on seed germination, seedlings growth and pigment content of Vigna mungo (L) Hepper (CVT 9)". *Geobios* 20 (1993): 108-112.
28. Kannan J. "Effect of distillery effluents on crop plants". *Advanced Plant Science* 14.1 (2001): 127-132.
29. Keraita B., *et al.* "On -farm treatment options for wastewater, greywater and fecal sludge with special reference to West Africa. Colombo, Sri Lanka: International Water Management Institute IWMI).CGIAR Research Program on Water, Land and Ecosystems (WLE)". Resource Recovery and Reuse Series (2014): 32.
30. Khurana MPS and Pritpal Singh. "Waste Water Use in Crop Production: A Review". *Resources and Environment* 2.4 (2012): 116-131.
31. Kumar Rajesh. "Effect of sugar mill effluent on the seed germination and seedling growth of Cicer arietinum". *Advanced Plant Science* 8.1 (1995): 25-26.
32. Mahapatro M and BK Mohanty. "Effect of Sugar Mills Effluents on germination of green gram (Phaseolus mungo Roxb.) and growth behaviour of its seedlings". *Jr. Indust. Polln. Contl* 23.1 (2007): 55-59.
33. Mahida UN. "Water Pollution and Disposal of Wastewater on Land". Tata McGraw-Hill & Co., New Delhi (1981):
34. Manivasakam N. "Physico Chemical Examination of Water, Sewage and Industrial Effluents". Pragati Prakashan, Meerut (2000):
35. Mathur A., *et al.* "A study of river Ganga at Varanasi with special emphasis on heavy metal pollution". *Pollution Research* 6.1 (1987): 37-44.
36. Mexico CAN. "Water Statistics; National Water Commission, Mexico City Muñoz". (2004):
37. Mishra V and SD Pandey. "Effect of distillery effluent and leachates of industrial sludge on the germination of Bengal gram (Cicer arietinum)". *Pollution Research* 21.4 (2002): 461-467.
38. Mishra SG., *et al.* "Relevance of sewage sludge in agriculture in 21st Century". *Vijana Parishad Anussandhan Patrika* 41.4 (1998):
39. Mitra AK. "Chemical characteristics of surface water at selected gauging stations in the river Godavari, Krishna and Tungabhadra at Kurnoll Town". *Indian Journal of Environmental Health* 24 (1982): 165-179.
40. Murthy GVR., *et al.* "A preliminary study on water quality of river Tungabhadra at Kurnoll Town". *Indian Journal of Environmental Health* 14.8 (1994): 604-607.
41. Muthukumar B and DI Arokiasamy. "Effects of distillery waste on the nitrate reduction system in Vigna mungo (Black gram)". *Journal of ecotoxicology & environmental monitoring* 4 (1994): 27-33.
42. Nagarajah S. "In: Green Manure in Rice Farming, IRRI, Philippines". (1988): 193.
43. Nauhbar Suman. "Relative tolerance of crop plants to salt stress at the early seedling stage; Ph.D. Thesis, Dr B R Ambedkar University, Agra". (2005):
44. NEERI. "Utilization of Treated Wastewaters from Mathura Refinery for Irrigation; NEERI, Nagpur". (1981):
45. NEERI. "Manual on Water and Waste Water Analysis; NEERI, Nagpur". (1988):
46. Parashar Nidhi. "Planning and Investigation for City and Industrial Effluent Utilization in Abating Pollution of River Yamuna and Improving Agricultural Production". Ph.D. Thesis, Dr B R Ambedkar University, Agra (2011):
47. Parashar Nidhi and Ravi Sharma. "Crop response to abiotic stresses of effluent, salinity and Heavy Metal: I – Comparative percent seed germination in mustard and wheat". *Agricultural Research & Technology* 3.3 (2017): 1-17.
48. Parashar Nidhi., *et al.* "City and Industrial Effluent Analysis at Five Selected Sites in Mathura UP India". *International Journal of Health, Safety and Environments* 4.1 (2018): 180-202.
49. Pathak Abha. "Differential growth and metabolic behaviour of cereal and legume crops under the influence of heavy metal cadmium". Ph.D. Thesis, Dr B R Ambedkar University, Agra (2017):
50. Pathak A., *et al.* "Differential Growth and Metabolic Behaviour of Cereal and Legume Crops under the Influence of Heavy Metal Cadmium: I – Germination Percentage (GP) and Speed of Germination Index (SGI)". *Journal of Plant Physiology* 1.1 (2017): 1-22.

51. Pedrero F, *et al.* "Use of treated municipal wastewater in irrigated agriculture. Review of some practices in Spain and Greece". *Agricultural Water Management* 97 (2010): 1233-1241.
52. Pragasam A and B Kannabiran. "Effect of distillery effluent on the morphology, biochemistry and yields of Vigna mungo". *Advanced Plant Science* 14.2 (2001): 547-552.
53. Pragasam A and B Kannabiran. "Effect of distillery effluent on the photosynthetic pigment and nitrogenous compound in Vigna mungo(L) Hepper". *J Swamy Bot* 18 (2001): 21-23.
54. Pragasam A and B Kannabiran. "Effect of distillery effluent on the NPK contents of Vigna mungo (L) Hepper and physico-chemical properties of soil". *Indian Journal of Environment and Ecoplanning* 8.2 (2004): 297-300.
55. Qadir M., *et al.* "The challenges of wastewater irrigation in developing countries". *Agricultural Water Management* 97.4 (2010): 561-568.
56. Rai H. "Limnological studies on the River Yamuna at Delhi, India, Part I: Relation between the chemistry and the state of pollution in the River Yamuna". *Archiv fur Hydrobiologie* 73 (1974): 369-393.
57. Rai H. "Limnological studies on the River Yamuna at Delhi, India, Part II: The dynamics of phytoplankton populations in the River Yamuna". *Archiv fur Hydrbiologie* 73 (1974): 492-517.
58. Raiyani CV, *et al.* "Magnitude of pollution of dye contaminated river water – Its physico-chemical and microbial analysis". *Indian Journal of Environmental Protection* 14.4 (1994): 252-255.
59. Rajendran K. "Effect of distillery waste on the seed germination, seedling growth, chlorophyll content and mitosis on Helianthus annus". *Indian Botanical Contracter* 7 (1990): 139-143.
60. Rani Saroj. "Investigations on salt tolerance parameters specially growth and biochemical traits for selection of salt tolerant lines in legumes at the early seedling stage; Ph.D. Thesis, Dr B R Ambedkar University, Agra". (2007):
61. Rani Saroj, *et al.* "Germination and early seedling growth in six leguminous pulse crops under salt stress". *Plant Archives* 9.1 (2009): 145-151.
62. Reid GK. "Ecology of Inland Waters and Estuaries; Reinhold Publishing Corp., New York". (1964):
63. Rusan MJM., *et al.* "Long term effect of waste water irrigation of forage crops on soil and plant quality parameters". *Desalinization* 215 (2007): 143-152.
64. Sahai R., *et al.* "Effect of distillery waste on seed germination, seedling growth arid pigment contents of rice". *Indian Ecological Society* 10 (1983): 7-10.
65. Sakai H. "A method for determining nitrifying power of paddy soil and some results with its use". *Progress Report of the All India Coordinated Rice Improvement Project 2, Kharif* (1970): 195.
66. Saraswat Kapila. "Studies on the phyto-toxicity of heavy metal mercury on germination, early seedling growth and biochemical changes in wheat (*Triticum aestivum* L)". Ph D Thesis, Dr B R Ambedkar Univ, Agra (2017):
67. Sarin MN and IM Rao. "Effect of sodium sulphate on early seedling growth of gram and wheat". *Agra Univ. J. Res* 5.1 (1956): 143-154.
68. Saxena KK and RS Chauhan. "Physico-chemical aspect of pollution at Agra". *Environmental Science and Pollution Research* 12.2 (1993): 101-104.
69. Sayato Y. "Hygienic chemical studies on effluent water, VI: Fundamental studies on hygienic chemistry of Brackish Lake". *International Journal of Hygiene and Environmental Health* 16 (1976): 32-38.
70. Sehgal JR and RH Siddiqui. "Characterization of wastewater from Kanpur City". *Environmental Health* 11 (1989): 95-107.
71. Shaji C and RJ Patel. "Chemical and biological assessment of pollution in the River Khari, Gujarat". *Indian Ecological Society* 17.1 (1990): 6-8.
72. Shani RL., *et al.* "Effect of distillery waste on seed germination, seedling growth and pigment contents of rice". *Indian Ecological Society* 10.1 (1983): 7-9.
73. Sharma Ajeet Kumar. "Water quality problem in Mathura and its effect on the physiology of major crops in the region". Ph. D. Thesis., Dr. B.R. Ambedkar Univ., Agra (2004):

74. Sharma AK, *et al.* "Study of Physico-chemical Characteristics of Water Quality of Yamuna River at Mathura, UP". *Bull. Env. Pharmacol. Life Sci* 2.10 (2013): 131-134.
75. Sharma AK, *et al.* "Monitoring of water quality of Yamuna River at Mathura, U.P. – Physico-Chemical Characteristics". *International Journal of Environmental Science and Technology* 3.4 (2013): 156-159.
76. Sharma Ajit Kumar, *et al.* "Impact of textile waste water on seed germination and growth development of *Vicia faba* L. at Jaipur, Rajasthan, India". *Journal of the Kalash Science* 2.2 (2014): 47-51.
77. Sharma, *et al.* "Impact of paper mill effluent on seed germination and seedling growth of *Cyamopsis tetragonoloba* L in Sanganer Region of Jaipur (Raj.) India". *Jour Environ. Sc., Computer Sc. And Engineer. & Tech* 3.4 (2014): 1830-1835.
78. Sharma, *et al.* "Impact of paper mill effluent on seed germination and seedling growth of *Vigna radiata* L. A case study in Sanganer Region at Jaipur (Raj.) India". *Cambridge Scholars Publishing, UK* 20 (2014): 1-5.
79. Sharma, *et al.* "Impact of Paper Mill Effluent on Seed Germination and Seedling Growth of *Vigna radiata* L.A Case Study in Sanganer Region at Jaipur (Raj) India". 20 (2016): 209-215.
80. Sharma AK, *et al.* "Physico-Chemical Characteristics of Yamuna River at Mathura District UP". *International Journal of Pharmaceutics* 7.1 (2018): 1-10.
81. Sharma HB, *et al.* "Water quality of sewage drains entering Yamuna River at Mathura (U.P)". *Journal of Environmental Biology* 21.4 (2000): 375-378.
82. Sharma Ravi. "Environmental Problems in Mathura; Workshop on Environmental Education for Asian Youth, INSA, New Delhi & Nainital (UP)". *Proc Indian Environmental Society* 56 (1976): 1.
83. Sharma Ravi. "Physiology of plant tolerance to salinity at early seedling stage". Ph D Thesis, Agra Univ, Agra (1982):
84. Sharma Ravi. "Towards an understanding of the physiology of salt tolerance in wheat (*Triticum aestivum* L.) at early seedling stage". XIV International Botanical Congress, Berlin (W), Germany 22a.8 (1987): 32.
85. Sharma Ravi. "Genotypic response to salt stress: I – Relative tolerance of certain wheat cultivars to salinity". *Advances in Crop Science and Technology* 3.4 (2015): 1-7.
86. Sharma, Ravi 2017 Genotypic response to salt stress: II – Pattern of Differential Relative Behaviour of Salt-tolerant, Moderately Salt-Tolerant and Salt-Sensitive Wheat cultivars under Salt Stressed Conditions; *Inter Jour Gr Sed & Water. (IJGSW), Univ. of Wisconsin, USA*, 5: 205 – 216.
87. Sharma V, *et al.* "Distillery effluent on seed germination, early seedling growth and pigment content of sugar beet". *Journal of Environmental Biology* 23.1 (2002): 77-80.
88. Sheoran IS and OP Garg. "Effect of salinity on the activities of RNase, DNase and protease during germination and early seedling growth of mung bean". *Plant physiology* 44.3 (1978): 171-174.
89. Shi R, *et al.* "Countermeasures of Reclaimed Municipal Wastewater for Safe ty of Agricultural Use in China". *Agricultural Sciences in China* 7.11 (2008): 1365-1373.
90. Singh DK. "Effect of heavy metal on productivity of grassland". Ph. D. Thesis., Banaras Hindu Univ, Varanasi (2003):
91. Singh DK, *et al.* "Physico-chemical analysis of effluents from a carpet industry". *Plant Archives* 8.2 (2008): 577-583.
92. Singh JPPK, *et al.* "Pollution status on Sangam and its adjoining river before the Kumbh Mela at Allahbad". *Indian Journal of Environmental Pollution* 8.11 (1988): 839-842.
93. Singh JPPK, *et al.* "Mass bathing effect on water quality of Sangam during Mahakumbh Mela at Allahbad". *Indian Journal of Environmental Pollution* 9.3 (1989): 189-193.
94. Singh JPPK, *et al.* "The assessment of water quality of Sangam and its adjoining river after the Mahakumbh Mela at Allahbad". *Indian Journal of Environmental Pollution* 9.5 (1989): 372-375.
95. Singh K.P, *et al.* "Impact assessment of treated/untreated wastewater toxicants discharge by sewage treatment plant on health, agricultural and environmental quality in waste water disposal area". *Chemosphere* 55.2 (2004): 227-255.
96. Singh Shalini. "Relative tolerance of crop plants to heavy metal pollution". Ph D Thesis, Dr B R Ambedkar Univ, Agra (2009):

97. Smith CJ and RD Delaune. "Effect of rice plants on nitrification, denitrification loss of nitrogen under greenhouse conditions". *Plant and Soil* 79.2 (1984): 287-290.
98. Snehlata. "Utilization of Sewage Effluents for Irrigating Crop Plants for Their Sustainable Improvement with special Reference to Mathura". Ph.D. Thesis, Dr. B R Ambedkar University, Agra (2016):
99. Srivastava RK. "Effects of distillery waste on the seed germination, early seedling growth and pigment content of Cajanas cajan L". *Indian Botanical Society* 65 (1986): 208-211.
100. Strauss M and UJ Blumenthal. "Human waste use in agriculture and aquiculture: utilization practices and health perspectives". IRCWD Report 09/90. International Reference Centre for waste Disposal 48 (1990):
101. Subramani A., et al. "Effect of distillery effluent on growth yield and productivity of Vigna radiata". *Pollution Research* 141 (1995): 477-482.
102. Sunderrajan KS., et al. "Biochemical monitoring of sewage pollution". *Proc. Acad. Environ. Biol* 3.1 (1994): 111-114.
103. Tebbutt THY. "Principles of Water Quality Control; Pergamon Press, London". (1977):
104. Thakur Raj Gopal. "Study of the pollutants of river Yamuna caused by the metallic Industries at Mathura". Ph. D. Thesis., Dr. B.R. Ambedkar Univ., Agra (2001):
105. Tiwari P., et al. "Responses of two cultivars of Indian mustard {[Brassica juncea (L) Zern & Coss]} to distillery". *Environmental Science and Pollution Research* 3.3 (1993): 131-143.
106. Tonapi GT. "Freshwater Animals of India; Oxford and IBH Pub. Co., New Delhi". (1980):
107. Trivedy RK and PK Goyal. "Chemical and Biological Methods for Water Pollution Studies". Environmental Publication, Karad, India (1986):
108. Unnamalai CT, et al. "Studies on the biofunctional efficiency of industrial and urban solid wastes on rice production based on their mineralization". *Plant Archives* 7.2 (2007): 705-707.
109. Weber CI. "A Guide to Common Diatoms of Water Pollution Surveillance System Stations; U.S.E.P.A. Cincinnati, Ohio". (1971):
110. WHO. "International Standards for Drinking Water; Geneva 3rd ed". (1984):
111. Yadav. "Physiology of salt tolerance for effective biological control of salinity". Ph D Thesis, Dr B R Ambedkar Univ, Agra (2006):
112. Yadav RS and RM Saxena. "Physico-chemical characterization of the tannery effluents of Agra". *Nature & Environment* 4 (1987): 50-52.

Submit your next manuscript to Scientia Ricerca Open Access and benefit from:

- Prompt and fair double blinded peer review from experts
- Fast and efficient online submission
- Timely updates about your manuscript status
- Sharing Option: Social Networking Enabled
- Open access: articles available free online
- Global attainment for your research

Submit your manuscript at:

<https://scientiaricerca.com/submit-manuscript.php>